

APTAMIL 3 PROFUTURA DUOBIOTIK

Date of issue: 01/03/2022 No.: 128853-001-002 Page 1/3

Powdered formula for babies in early infancy (from 12 months) based on skimmed milk

From month

Our unique combination:

- HMO* (2'FL and 3'GL**)
- Prebiotic fibres (GOS/FOS)
- Milk fat
- OMEGA-3
- Immunonutrients (vitamins C and D)
- Football

AVAILABLE FORMATS

EZTin: 800 g

SHELF LIFE: 18 months

RECONSTITUTION

The reconstitution of Aptamil®Profutura® 3 Duobiotik is 15.1% and is achieved by adding 1 level scoop of powder (approx. 5 g) to every 30 ml of water.

METHODS OF STORAGE AND USE

WARNING: Follow the instructions carefully: improper preparation and storage may pose a risk to health. health of the infant.

Before preparation, make sure the working environment is clean and wash and dry your hands well.

Sterilise the bottle, nipple and ring. If you do not have a steriliser, boil them in water.

Boil the amount of drinking water required for reconstituting the product in a clean container and allow it to cool.

Pour lukewarm water into the sterilised feeding bottle according to the instructions in the media feeding table.

Add the level scoops of powder according to the indications on the average feeding chart using only the scoop contained in the pack.

Close the bottle and shake it for at least 10 seconds to dissolve the powder.

Once the product is ready, test the temperature on the pulse and administer to the infant. Consume within half an hour of preparation.

Do not reuse any leftover prepared product, and do not heat or boil milk once it has been reconstituted.

After use, close the package tightly and store in a clean, cool, dry place (not in the refrigerator). Consume within 1 month of opening.

PACKAGE DESIGNED FOR YOU

Triple security seal: to make sure the package has not been opened

Safety tab: for greater safety against contamination and spillage

Measuring cup holder: to keep the measuring cup and hands away from dust

AVERAGE SUPPLY TABLE

The table is indicative, the dosage may be modified according to the advice of the paediatrician, depending on the infant's needs. Use only the measuring cup contained in the pack.

| i . | | | 1 |
|-------------|---------------------|-----------------|---|
| To prepare | No. of level scoops | Boiled water ml | |
| one cup | 7 | 210 | |

IMPORTANT WARNINGS

Breast milk is the ideal food for the baby and is recommended as the exclusive food until the sixth month and as the milk component of a balanced diet during weaning and also after one year of life. Aptamil®Profutura® 3 Duobiotik is intended as part of a varied and balanced diet and a healthy lifestyle, following the paediatrician's instructions.

NUTRITIONAL CHARACTERISTICS

Our best Aptamil® Profutura® 3 Duobiotik formula inspired by the latest research on breast milk. Our unique combination of:

HMO 2'FL, HMO 3'GL* and Prebiotic Fibres (GOS/FOS).

Milk fat

Immunonutrients: vitamins C and D contribute to the normal functioning of the immune system

Omega-3 (ALA and DHA)*: ALA contributes to the development of the brain and nerve tissue

Calcium: contributes to normal bone development

3'GL resulting from our unique process

Product packed in a protective atmosphere.

Distributed by: Danone Nutricia S.p.A. Società Benefit, Via Carlo Farini 41, 20159 Milan (MI).

Best before the date on the pack. Store in a cool, dry place, away from heat sources.

APTAMIL 3 PROFUTURA DUOBIOTIK

Date of issue: 01/03/2022 No.: 128853-001-002 Page 2/3

INGREDIENTS

Lactose, skimmed milk powder, maltodextrins, galacto-oligosaccharides from milk (powder), whey products powder (demineralised whey, whey protein concentrate), rapeseed oil low in erucic acid, sunflower oil, coconut oil, anhydrous milk fat, calcium phosphate, sunflower oil high in oleic acid, fructooligosaccharides, potassium hydrogen phosphate, potassium citrate, fish oil, magnesium chloride, 2'-fucosyl lactose, sodium chloride, L-ascorbic acid, magnesium hydrogen phosphate, sodium citrate, choline chloride, emulsifier: soy lecithin; ferrous sulphate, potassium chloride, inositol, sodium L-ascorbate, antioxidant: acorbyl palmitate; zinc sulphate, L- carnitine, calcium D-pantothenate, DL-α-tocopheryl acetate, calcium carbonate, nicotinamide, riboflavin, retinyl palmitate, pyridoxine hydrochloride, thiamine hydrochloride, potassium iodide, folic acid, phytomenadione, cholecalcferol, D-biotin, cyanocobalamin.

| AVERAGE NUTRITIONAL VALUES | | per 100 g of dust | per 100 ml at | %AR * |
|---------------------------------|-------|----------------------|------------------|-------|
| AVERAGE NOTRITIONAL VALUES | | or dust | 15.1% | %AK " |
| Energy | kJ | 73 | 18112 | |
| | kcal | | 43165 | |
| Fats | g | 6 | 17,22, | |
| of which saturated fatty acids | g | | 520,79 | |
| Carbohydrates | g | 7 | 57,88, | |
| of which Sugars | g | 9 | 45,96, | |
| Fibres | g | 9 | 580,9 | |
| Protein | g | | 861,3 | |
| Halls | g | 06 | 0,430, | |
| Vitamins and minerals | | | | |
| Vitamin D | μg | | 213,1 | 44% |
| Vitamin E | mg TE | | 6,91,0 | 20% |
| Vitamin K | μg | | 345,1 | 43% |
| Vitamin C | mg | | 9915 | 33% |
| Riboflavin (vitamin B2) | mg | | 150,23 | 33% |
| Vitamin B12 | μg | | 2,70,4 | 50% |
| Pantothenic acid | mg | 583 | 3,8670, | 19% |
| Biotin | μg | | 111,6 | 16% |
| Football | mg | 0 | 79612 | 22% |
| Phosphorus | mg | | 63796 | 18% |
| Potassium | mg | 0 | 99815 | 15% |
| Iron | mg | ٧A). | 821,2 | 15% |
| lodine | μg | 7 | 13220 | 25% |
| | // | | | |
| Other nutrients | NA | | O/X | |
| linoleic acid (LA) | mg | 52 | 37005 | |
| α-linolenic acid (ALA) | mg | 4 | 65098, | |
| eicosapentaenoic acid (EPA) | mg | - 4 | 7010,4 | |
| docosahexaenoic acid (DHA) | mg | 5 | 10015, | |
| lactose | g | 9 | 4246,3 | |
| GOS (non-caloric carbohydrates) | g | - | 230,3 | |
| 2'-Fucosyl Lactose (2'FL) | g | 02 | 0,130, | |
| | | | | |

^{*} Reference assumptions Delegated Regulation (EU) 2016/127

APTAMIL 3 PROFUTURA DUOBIOTIK

Date of issue: 01/03/2022 No.: 128853-001-002 Page 3/3

SUBSTANCES AND PRODUCTS CAUSING ALLERGIES OR INTOLERANCES

| (EU Reg. 1169/2011 Annex 2) | | |
|--|----|----|
| <u>></u> | SI | NO |
| 1. Cereals containing gluten, namely: wheat (spelt and khorasan wheat), rye, barley, oats or their (a) wheat-based glucose syrups, including dextrose (1); (b) wheat-based maltodextrins (1); (c) glucose syrups based on barley; (d) cereals used for making alcoholic distillates, including ethyl alcohol of agricultural origin. | | х |
| 2. Crustaceans and crustacean products. | | X |
| 3. Eggs and egg products. | | Χ |
| Fish and fish products, except: (a) fish gelatine used as a carrier for vitamin or carotenoid preparations; (b) gelatine or isinglass used as a fining agent in beer and wine. | х | |
| 5. Peanuts and peanut products. | | X |
| 6. Soy and soy products, except: (a) refined soybean oil and fat (1); (b) natural mixed tocopherols (E306), natural D-alpha tocopherol, natural D-alpha tocopherol acetate, (c) vegetable oils derived from phytosterols and phytosterol esters based on soy; (d) Plant stanol ester produced from soybean-based vegetable oil sterols. | х | |
| 7. Milk and milk products (including lactose), except:(a) whey used in the manufacture of alcoholic distillates, including ethyl alcohol of agricultural origin;(b) lactiol. | x | |
| 8. Nuts, i.e.: almonds (Amygdalus communis L.), hazelnuts (Corylus avellana), walnuts (Juglans regia), cashews (Anacardium occidentale), pecans (Carya illinoinensis (Wangenh.) K. Koch], Brazil nuts (Bertholletia excelsa), pistachio nuts (Pistacia vera), macadamia nuts or Queensland nuts (Macadamia ternifolia), and their products, except for nuts used for the manufacture of alcoholic distillates, including ethyl alcohol of agricultural origin. | | х |
| 9. Celery and celery products. | | Х |
| 10. Mustard and mustard products. | | Х |
| 11. Sesame seeds and sesame seed products. | | Х |
| 12. Sulphur dioxide and sulphites at concentrations of more than 10 mg/kg or 10 mg/litre in terms of total SO 2 to be calculated for the products as proposed ready for consumption or as reconstituted according to the instructions of the manufacturers. | | х |
| 13. Lupins and lupin products. | | Х |
| 14. Molluscs and mollusc products. | | Х |

⁽¹⁾ And derived products, insofar as the processing which they have undergone is not likely to increase the level of allergenicity assessed by the Authority for the commodity from which they are derived.

REFERENCE LEGISLATION

Reg. 1925/2006 "On the addition of vitamins and minerals and of certain other substances to foods" and Ministry of Health Note of 05/07/2016.

Delegated Regulation (EU) 2016/127. Supplements Regulation (EU) No 609/2013 of the European Parliament and of the Council as regards specific compositional and information requirements for infant formulae and follow-on formulae and as regards information requirements for infant and young child feeding.